



**FIG. 3.** Image shows the nasal septum after excision of the mass. NS: Nasal septum, IT: Inferior turbinate.



**FIG. 4.** A high-power view a lobule shows the compact proliferative of capillaries (H&E x400).

LCH which is also known as pyogenic granuloma typically occurs on the skin and in the oral cavity. Nasal cavity is an uncommon area for LCH (1). It is most frequently seen in the third and fifth decades of life, especially most commonly in women. Trauma and hormonal factors are considered for the etiology (2). LCH is usually violaceous, ulcerous, pedunculated or sessile lesions which bleeds on touch. It has variable sizes



**FIG. 5.** A low-power view a lobule shows the compact proliferative of capillaries (H&E x40).

ranging from a few millimeters to a few centimeters. Nonspecific symptoms such as epistaxis, nasal obstruction, and purulent rhinorrhea have been reported in majority of the patients (3). CT or magnetic resonance imaging is useful for preoperative evaluation. CT is important for assessing of nasal and paranasal bone structures, particularly for the large-sized lesions which is originate from the nasal roof because the osseous destruction of the skull base can be seen (4). Endoscopic surgery is the preferred approach for the treatment (5).

**Conflict of Interest:** No conflict of interest was declared by the authors.

## REFERENCES

1. Derkenne R, Coulet O, Varoquaux A, de Biasi C, Tomasi M. Nasal cavity lobular capillary hemangioma due to insect sting. *Eur Ann Otorhinolaryngol Head Neck Dis.* 2012;129:278-80.
2. Chi TH, Yuan CH, Chien ST. Lobular capillary hemangioma of the nasal cavity: a retrospective study of 15 cases in taiwan. *Balkan Med J.* 2014;31:69-71.
3. Ozcan C, Apa DD, Görür K. Pediatric lobular capillary hemangioma of the nasal cavity. *Eur Arch Otorhinolaryngol.* 2004;261:449-51.
4. Karagama YG, Howarth K, Steel PR, Spencer MG. Lobular capillary haemangioma of the nasal vestibule: a rare entity. *Int J Pediatr Otorhinolaryngol.* 2002;66:71-5.
5. Hanazawa T, Yonekura S, Nakamura H, Fujikawa A, Okamoto Y. Pre-operative effects of the administration of systemic corticosteroids combined with antibiotics on a lobular capillary hemangioma in the nasal cavity. *Auris Nasus Larynx.* 2016;43:203-6.