

Not All Brugada Electrocardiogram Patterns are Brugada Syndrome or Brugada Phenocopy

Grace Xu¹, Byron H. Gottschalk², Umut Kocabaş³, Adrian Baranchuk¹

¹Department of Medicine, Queen's University School of Medicine, Kingston, Canada

²Department of Cardiac Surgery, Western University School of Medicine, Ontario, Canada

³Clinic of Cardiology, Soma State Hospital, Manisa, Turkey

We have recently read the case report by Arı and Ekici (1) with great interest and noted that this patient displayed electrocardiogram patterns characteristic of Brugada syndrome after a propafenone overdose with the intention of suicide. This case report is an important contribution to the investigation of the aetiology of Brugada electrocardiogram patterns and their classification as Brugada syndrome, Brugada phenocopies, or other diagnoses. We have published extensively on the topic of Brugada phenocopies (2,3) and would like to clarify the terminology used in classifying the case report to avoid confusion in future research.

Antiarrhythmic agents block sodium channels and may be used to unmask true Brugada syndrome at appropriate dosages (4). The class 1C antiarrhythmic agent, propafenone, taken at equivalent doses, may serve a similar role to other sodium channel blockers but has not been systematically tested. Although the effects of large doses have not been systematically investigated, we suspect that overdoses of propafenone may induce artificial sodium channel dysfunction even without genetic predisposition, causing a Brugada electrocardiogram pattern.

The term Brugada phenocopies does not cover abnormal electrocardiogram patterns that are induced by sodium channel blockers (2). Thus, overdose using propafenone would not be considered Brugada phenocopies and instead, may be classified as an "acquired sodium channel dysfunction".

We thank Dr. Arı and Ekici for using the term Brugada phenocopies in their literature; however, we would like to emphasise that Brugada electrocardiogram patterns induced by sodium channel blockers should not be classified as Brugada phenocopies. We are continuing the development of our international online registry at www.brugadaphenocopy.com and encourage investigators working on Brugada phenocopies to submit their cases online to provide long-term follow up and investigation of the natural history of this condition.

Conflict of Interest: No conflict of interest was declared by the authors.

REFERENCES

1. Arı ME, Ekici F. Brugada-Phenocopy Induced by Propafenone Overdose and Successful Treatment: A Case Report. *Balkan Med J* 2017;5:458-63.
2. Baranchuk A, Nguyen T, Ryu MH, Femenia F, Zareba W, Wilde AA, et al. Brugada phenocopy: new terminology and proposed classification. *Ann Noninvasive Electrocardiol* 2012;17: 299-314.
3. Anselm DD, Baranchuk A. Brugada phenocopy: redefinition and updated classification. *Am J Cardiol* 2013;111:453.
4. Antzelevitch C, Brugada P, Borggrefe M, Brugada J, Brugada R, Corrado D, et al. Brugada syndrome: report of the second consensus conference: endorsed by the Heart Rhythm Society and the European Heart Rhythm Association. *Circulation* 2005;111:659-70.

Address for Correspondence: Dr. Grace Xu, Department of Medicine, Queen's University School of Medicine, Kingston, Canada

Phone: 647 234 6926

e-mail: 15gjx@queensu.ca

Received: 31 May 2017

Accepted: 15 September 2017 • DOI: 10.4274/balkanmedj.2017.0621

Available at www.balkanmedicaljournal.org

Cite this article as:

Xu G, Gottschalk BH, Kocabaş U, Baranchuk A. Not All Brugada Electrocardiogram Patterns are Brugada Syndrome or Brugada Phenocopy. *Balkan Med J* 2017;34:593

©Copyright 2017 by Trakya University Faculty of Medicine / The Balkan Medical Journal published by Galenos Publishing House.

