



Overlap Between Nutritional Indices in Patients with Acute Coronary Syndrome: A Focus on Albumin

✉ Mert İlker Hayiroğlu¹, ✉ Servet Altay²

¹Department of Cardiology, University of Health Sciences Türkiye, Dr. Siyami Ersek Thoracic and Cardiovascular Surgery Training and Research Hospital, İstanbul, Türkiye

²Department of Cardiology, Trakya University Faculty of Medicine, Edirne, Türkiye

Albumin, a key protein synthesized in the liver, plays a crucial role beyond its traditional function of maintaining oncotic pressure and fluid balance. Recently, its use as a prognostic marker for cardiovascular diseases (CVDs) has garnered attention. Serum albumin levels below normal have been repeatedly linked to higher rates of morbidity and mortality in individuals with heart diseases, particularly those with acute coronary syndrome (ACS).¹ This association highlights albumin's potential utility as a biomarker for risk stratification and mortality prediction in CVD. Understanding the mechanisms linking albumin deficiency to adverse cardiovascular outcomes may offer insights into novel therapeutic approaches and personalized management strategies. This editorial explores the current evidence on the predictive value of albumin in CVD and discusses its implications for clinical practice and future research directions.

The role of albumin as a prognostic indicator in patients with ACS has gained acceptance following the demonstration of its predictive value for both in-hospital and long-term mortality outcomes in multiple studies.² Therefore, numerous indices incorporating the albumin level have been established. Furthermore, several additional parameters have been added to the albumin level to better reach the endpoints. The prognostic nutritional index, which was first established for patients with cancer, was initially applied in cardiology in patients with ACS.^{3,4} Subsequently, the geriatric nutritional index and controlling nutritional (CONUT) score have been found to influence significant outcomes in both acute and chronic CVDs. Similar to studies on the prognostic nutritional index, the most frequently studied disease using these indices is ACS.^{5,6} The total cholesterol level, lymphocyte count, and weight of the patients were added to these indices to increase their acuity in predicting both in-hospital and long-term outcomes. In the prognostic nutritional index, lymphocyte count was integrated with the albumin level. Similarly, the CONUT score incorporates both the lymphocyte count and total cholesterol level. In the geriatric nutritional index, patient weight

has been included with the albumin level (Figure 1). These scores represent significant examples illustrating the value of albumin in patients with ACS. From this perspective, the crucial consideration arises: why does albumin impact prognostication and consistently demonstrate a significant relevance across multiple studies?

Several important physiological and clinical systems must be considered to determine the possible causes of albumin's predictive role. First, the albumin level reflects a patient's systemic inflammation and general nutritional state. Low albumin levels are frequently associated with chronic inflammation, malnourishment, and compromised cellular repair mechanisms. These factors play a crucial role in the development of adverse events and the course of coronary artery disease.⁷ The pathophysiology of coronary artery disease is dependent on endothelial dysfunction, plaque instability,




FIG. 1. The inclusion of the patient's weight into the albumin level in the geriatric nutritional index.

Corresponding author: Mert İlker Hayiroğlu, Department of Cardiology, University of Health Sciences Türkiye, Dr. Siyami Ersek Thoracic and Cardiovascular Surgery Training and Research Hospital, İstanbul, Türkiye

e-mail: mertilkerh@gmail.com

Available Online Date: xxxxx • **DOI:** 10.4274/balkanmedj.galenos.2024.190724

Available at www.balkanmedicaljournal.org

ORCID iDs of the authors: M.I.H. 0000-0001-6515-7349; S.A. 0000-0001-7112-3970.

Cite this article as: Hayiroğlu Mİ, Altay S. Overlap Between Nutritional Indices in Patients with Acute Coronary Syndrome: A Focus on Albumin. *Balkan Med J.*

Copyright@Author(s) - Available online at <http://balkanmedicaljournal.org/>

and increased susceptibility to thrombosis, which are exacerbated by malnutrition and inflammation. Furthermore, albumin acts as a store for bioactive compounds such as antioxidants and binding proteins, which include medications and hormones. Its antioxidant properties help protect against oxidative stress, a major contributor to endothelial dysfunction and atherosclerosis progression. Low albumin levels can cause a reduction in antioxidant capacity, which aggravates the oxidative damage to the arterial wall and encourages the development of atherogenesis.⁸ Albumin also regulates endothelial function and vascular tone. It modulates nitric oxide synthesis, a key vasodilator that maintains vascular homeostasis. Low albumin levels can impair nitric oxide bioavailability, leading to vasoconstriction, increased vascular resistance, and compromised coronary perfusion, which predispose patients to ischemic events.⁹ Beyond its direct physiological roles, albumin serves as a surrogate marker for systemic comorbidities and severity of coronary artery disease. Low albumin levels are associated with a higher burden of cardiovascular risk factors such as hypertension, diabetes mellitus, and chronic kidney disease, all of which independently contribute to adverse cardiovascular outcomes.¹⁰ Furthermore, albumin levels may reflect the presence of acute stressors and acute-phase reactions, which are common in patients experiencing adverse cardiac events. Albumin's predictive value in coronary artery disease extends to its association with clinical outcomes such as myocardial infarction, stroke, and cardiovascular mortality. Studies have consistently demonstrated that low serum albumin levels independently predict the incidence of adverse cardiac events and are associated with a poor prognosis in patients with coronary artery disease who are undergoing coronary interventions or medical therapy.¹¹ This prognostic utility highlights albumin's role as a valuable biomarker for risk stratification and clinical management in coronary artery disease. The multifaceted role of albumin in predicting adverse cardiac events in patients with coronary artery disease can be attributed to its reflection of nutritional status, inflammatory burden, antioxidant capacity, vascular function, and overall disease severity.¹² Furthermore, its predictive power highlights the complex interplay between systemic inflammation, oxidative stress, endothelial dysfunction, and atherosclerosis progression, which contributes to the pathophysiology of coronary artery disease. However, further studies are required to elucidate the precise mechanisms linking albumin level to adverse cardiovascular outcomes and explore albumin's potential as a therapeutic target in mitigating cardiovascular risk in patients with coronary artery disease.

In conclusion, the potential for including albumin levels in prediction models for both in-hospital and long-term mortality associated with CVD, particularly ACS, is significant. In the realm of personalized medicine, machine learning models and artificial

intelligence systems are poised to play central roles in future advancements in coronary artery disease.¹³ Furthermore, albumin has been integral in constructing predictive nomograms for both cardiovascular and non-CVD.^{14,15}

Authorship Contributions: Concept- M.I.H., S.A.; Design- M.I.H., S.A.; Data Collection or Processing- M.I.H., S.A.; Analysis or Interpretation- M.I.H., S.A.; Literature Search- M.I.H., S.A.; Writing- M.I.H., S.A.

Conflict of Interest: No conflict of interest was declared by the authors.

REFERENCES

- Zhu L, Chen M, Lin X. Serum albumin level for prediction of all-cause mortality in acute coronary syndrome patients: a meta-analysis. *Biosci Rep*. 2020;40:BSR20190881. [\[CrossRef\]](#)
- Manolis AA, Manolis TA, Melita H, Mikhailidis DP, Manolis AS. Low serum albumin: A neglected predictor in patients with cardiovascular disease. *Eur J Intern Med*. 2022;102:24-39. [\[CrossRef\]](#)
- Keskin M, Hayiroğlu MI, Keskin T, et al. A novel and useful predictive indicator of prognosis in ST-segment elevation myocardial infarction, the prognostic nutritional index. *Nutr Metab Cardiovasc Dis*. 2017;27:438-446. [\[CrossRef\]](#)
- Nozoe T, Ninomiya M, Maeda T, Matsukuma A, Nakashima H, Ezaki T. Prognostic nutritional index: a tool to predict the biological aggressiveness of gastric carcinoma. *Surg Today*. 2010;40:440-3. [\[CrossRef\]](#)
- Li Y, Shen J, Hou X, et al. Geriatric nutritional risk index predicts all-cause mortality in the oldest-old patients with acute coronary syndrome: A 10-year cohort study. *Front Nutr*. 2023;10:1129978. [\[CrossRef\]](#)
- Takahashi T, Watanabe T, Otaki Y, et al. Prognostic significance of the controlling nutritional (CONUT) score in patients with acute coronary syndrome. *Heart Vessels*. 2021;36:1109-1116. [\[CrossRef\]](#)
- Kurtul A, Murat SN, Yarlioglu M, et al. Usefulness of Serum Albumin Concentration to Predict High Coronary SYNTAX Score and In-Hospital Mortality in Patients With Acute Coronary Syndrome. *Angiology*. 2016;67:34-40. [\[CrossRef\]](#)
- Sastre-Oliva T, Corbacho-Alonso N, Rodriguez-Sanchez E, et al. Albumin Redox Modifications Promote Cell Calcification Reflecting the Impact of Oxidative Status on Aortic Valve Disease and Atherosclerosis. *Antioxidants (Basel)*. 2024;13:108. [\[CrossRef\]](#)
- Hattori Y, Banba N, Gross SS, Kasai K. Glycated serum albumin-induced nitric oxide production in vascular smooth muscle cells by nuclear factor kappaB-dependent transcriptional activation of inducible nitric oxide synthase. *Biochem Biophys Res Commun*. 1999;259:128-132. [\[CrossRef\]](#)
- Brobak KM, Halvorsen LV, Aass HCD, et al. Novel biomarkers in patients with uncontrolled hypertension with and without kidney damage. *Blood Press*. 2024;33:2323980. [\[CrossRef\]](#)
- Li M, Tang C, Luo E, Qin Y, Wang D, Yan G. Relation of Fibrinogen-to-Albumin Ratio to Severity of Coronary Artery Disease and Long-Term Prognosis in Patients with Non-ST Elevation Acute Coronary Syndrome. *Biomed Res Int*. 2020;2020:1860268. [\[CrossRef\]](#)
- Artigas A, Wernerman J, Arroyo V, Vincent JL, Levy M. Role of albumin in diseases associated with severe systemic inflammation: Pathophysiologic and clinical evidence in sepsis and in decompensated cirrhosis. *J Crit Care*. 2016;33:62-70. [\[CrossRef\]](#)
- Hayiroğlu MI, Altay S. The Role of Artificial Intelligence in Coronary Artery Disease and Atrial Fibrillation. *Balkan Med J*. 2023;40:151-152. [\[CrossRef\]](#)
- Zhou J, Lee S, Liu Y, et al. Predicting Stroke and Mortality in Mitral Regurgitation: A Machine Learning Approach. *Curr Probl Cardiol*. 2023;48:101464. [\[CrossRef\]](#)
- Gündoğan Uzunay B, Köker A, Ülgen Tekerek N, Dönmez L, Dursun O. Role of Albumin-corrected Anion Gap and Lactate Clearance in Predicting Mortality in Pediatric Intensive Care Patients. *Balkan Med J*. 2023;40:430-434. [\[CrossRef\]](#)