



OPEN ACCESS

In Regard to Bozkurt et al.

To the Editor,

I read the article by Bozkurt and Şahin (1) with great interest. I have some points of concern related to this case report.

First, the authors have stated that painful erections lasted for about 1 hours after adding olanzapine 2.5 mg/day. However, there was no comment regarding the number of attacks or the time of attacks. I believe that it is essential to determine whether periods of priapism occur during the day or the night in a 9-year-old boy because of possible physiological nocturnal erections in the peripubertal interval.

Second, in this case, priapism attacks were not observed by a health professional. Pain during erections or the color of the penis may lead us to a diagnosis of priapism; however, I believe that it should be clarified by an observation in the hospital. In addition, there are case reports of priapism with methylphenidate treatment (2,3). It would be safer to observe the patient for a few hours after the initiation of methylphenidate in case of recurrence of priapism.

Finally, the authors have stated that other organic causes of priapism were excluded. However, the details of the examination were not mentioned in the article. I believe that the details of any radiological

evaluation performed should be mentioned. This is because the diagnosis and treatment of priapism are not very clear for this age interval and there are only a few guidelines in the pediatric urology literature (4).

Mete Özkıdık

Clinic of Urology, Şanlıurfa Training and Research Hospital, Şanlıurfa, Turkey

Anar İbrahimov

Department of Urology, Ankara University School of Medicine, İbni Sina Hospital, Ankara, Turkey

REFERENCES

1. Bozkurt H, Şahin S. Olanzapine-induced Priapism in a Child with Asperger's Syndrome. *Balkan Med J* 2017;34:85-7.
2. Kelly BD, Lundon DJ, McGuinness D, Brady CM. Methylphenidate-induced erections in a prepubertal child. *J Pediatr Urol* 2013;9:1-2.
3. Esnafoglu E, Demir EY. Hallucination and priapism associated with methylphenidate usage: Two case reports. *Indian J Psychiatry* 2017;59:386-7.
4. Donaldson JF, Rees RW, Steinbrecher HA. Priapism in children: a comprehensive review and clinical guideline. *J Pediatr Urol* 2014;10:11-24.

Address for Correspondence: Dr. Mete Özkıdık, Clinic of Urology, Şanlıurfa Training and Research Hospital, Şanlıurfa, Turkey

e-mail: meteozkdk@gmail.com ORCID ID: orcid.org/0000-0002-7304-9396

Received: 26 February 2018 Accepted: 9 April 2018 • DOI: 10.4274/balkanmedj.2018.0407

Available at www.balkanmedicaljournal.org

Cite this article as:

Özkıdık M, İbrahimov A. In Regard to Bozkurt et al. *Balkan Med J* 2018;35:354

©Copyright 2018 by Trakya University Faculty of Medicine / The Balkan Medical Journal published by Galenos Publishing House.



In Reply to Özkıdık et al.

To the Editor,

We read the letter that criticizes our previous report published in *Balkan Medical Journal*. Commentaries to the letter are as noted below respectively (1).

Firstly, information regarding the medical and psychiatric history of this child case was taken from mother and we unfortunately did not have the chance to observe the penile erection attacks in a clinical setting. So we reported this pediatric case based on the mother's subjective observations. Those painless erections were only associated with olanzapine but not with any other factors or periods like day or night and resolved spontaneously after ceasing the

treatment. According to mother, her son has never experienced such like these penile erections leading her to cease the drug. However we agree that information about the number of attacks should have been indicated in the case report.

Secondly, we already stated that methylphenidate does not seem to cause priapism since the case benefited from methylphenidate without any side effect except loss of appetite. Naranjo algorithm for olanzapine rather than methylphenidate also classed the adverse reaction as "most probably" (1). Additionally, the case kept using methylphenidate while priapism resolved after ceasing olanzapine.

Address for Correspondence: Dr. Hasan Bozkurt, Department of Child and Adolescent Psychiatry, Gaziosmanpaşa University Hospital, Tokat, Turkey

e-mail: drhasan33@gmail.com

DOI: 10.4274/balkanmedj.2018.1180

Available at www.balkanmedicaljournal.org

Cite this article as:

Bozkurt H, Şahin S. In Reply to Özkıdık et al. *Balkan Med J* 2018;35:354-355

©Copyright 2018 by Trakya University Faculty of Medicine / The Balkan Medical Journal published by Galenos Publishing House.

